arXiv:hep-ph/9812247v2 14 Dec 1998

Physics with $B \to \eta'$ and $B \to J/\psi$ [a]

Xiao-Gang He
Department of Physics, National Taiwan University, Taipei, 10764, Taiwan

In this talk I discuss two topics related to B decays. I will first discuss the Standard Model prediction for $B \to \eta' X_s$, and then discuss how to use experimental information from $B \to J/\psi K^*$ to test CP violation beyond the SM.

I will report some of my recent work on $B \to \eta' X_s$ and $B \to J/\psi K^*$ with my collaborators [1,2,3] in this talk. The processes $B \to \eta' X_s$ and $B \to J/\psi K^*$ have been analyzed in detail by the CLEO collaboration [4,5]. The unexpected large branching ratio of [4] $(6.2 \pm 1.6 \pm 1.3) \times 10^{-4}$ for $B \to \eta' X_s$ has led some to claim that new physics is needed to explain the observation by CLEO [6,7,8]. I will show that in fact the Standard Model prediction is consistent with experimental data when related processes are treated properly [1]. The measurement of the full angular analysis of [5] $B \to J/\psi K^*$ from CLEO has also led to some interesting physics. I will show that information obtained from the angular distribution for this decay provide good test for CP violation beyond the SM [2,3].

1 $B \to \eta' X_s$ in the Standard Model

The recent observation of $B \to \eta' X_s$ [4] and $B \to \eta' K$ [9] decays with high momentum η' has stimulated many theoretical activities [6,7,8,10,11,12,13,14]. One of the mechanisms proposed to account for this decay is $b \to s g^* \to s g \eta'$ [6,10] where the η' meson is produced via the anomalous $\eta' - g - g$ coupling. According to a previous analysis [6], this mechanism within the Standard Model(SM) can only account for 1/3 of the measured branching ratio [4]. There are also other calculations of $B \to \eta' X_s$ based on four-quark operators of the effective weak-Hamiltonian [7,11]. These contributions to the branching ratio, typically 10^{-4}, are also too small to account for $B \to \eta' X_s$, although the four-quark-operator contribution is capable of explaining the branching ratio for the exclusive $B \to \eta' K$ decays [12,13]. These results have inspired proposals for an enhanced $b \to sg$ and other mechanisms arising from physics beyond the SM [6,7,8]. It will be shown here that the SM is in fact consistent with experimental data from CLEO [1].

---

[a] Talk presented at the 4th International Workshop on Particle Physics Phenomenology, Kaohsiung, Taiwan, China, 18-21 June, 1998

The quark level effective Hamiltonian for the $B \to \eta' X_s$ decay is given by[15]:

$$\begin{aligned} H_{eff}(\Delta B = 1) &= \frac{G_F}{\sqrt{2}}[\sum_{f=u,c} V_{fb}V^*_{fs}(C_1(\mu)O^f_1(\mu) + C_2(\mu)O^f_2(\mu)) \\ &- V^*_{ts}V_{tb}\sum_{i=3}^{6}(C^{eff}_i(\mu)O_i(\mu) + C^{eff}_8(\mu)O_8(\mu))], \qquad (1) \end{aligned}$$

The operators are defined in Ref.[15]. C_i are the Wilson Coefficients (WC). The values for the WC's given in Ref.[16,17] will be used in this discussion. The superscript ^{eff} indicate that the matrix element corrections are also included.

Let me first discuss the four-quark operator contributions to $B \to \eta' X_s$. The four-quark operators can induce three types of processes represented by 1) $< \eta'|\bar{q}\Gamma_1 b|B >< X_s|\bar{s}\Gamma'_1 q|0 >$, 2) $< \eta'|\bar{q}\Gamma_2 q|0 >< X_s|\bar{s}\Gamma b|B >$, and 3) $< \eta' X_s|\bar{s}\Gamma_3 q|0 >< 0|\bar{q}\Gamma'_3|B >$. Here $\Gamma^{(')}_i$ denotes appropriate gamma matrices. The contribution from 1) gives a "three-body" type of decay, $B \to \eta' s\bar{q}$. The contribution from 2) gives a "two-body" type of decay $b \to s\eta'$. And the contribution from 3) is the annihilation type which is relatively suppressed and will be neglected. Several decay constants and form factors needed in the calculations are listed below:

$$\begin{aligned} &< 0|\bar{u}\gamma_\mu\gamma_5 u|\eta' >=< 0|\bar{d}\gamma_\mu\gamma_5 d|\eta' >= i f^u_{\eta'} p^{\eta'}_\mu \\ &< 0|\bar{s}\gamma_\mu\gamma_5 s|\eta' >= i f^s_{\eta'} p^{\eta'}_\mu, \;\; < 0|\bar{s}\gamma_5 s|\eta' >= i(f^u_{\eta'} - f^s_{\eta'})\frac{m^2_{\eta'}}{2m_s}, \\ &f^u_{\eta'} = \frac{1}{\sqrt{3}}(f_1\cos\theta_1 + \frac{1}{\sqrt{2}}f_8\sin\theta_8), \;\; f^s_{\eta'} = \frac{1}{\sqrt{3}}(f_1\cos\theta_1 - \sqrt{2}f_8\sin\theta_8), \\ &< \eta'|\bar{q}\gamma_\mu b|B >= F^{Bq}_1(p^B_\mu + p^{\eta'}_\mu) + (F^{Bq}_0 - F^{Bq}_1)\frac{mB^2 - m^2_{\eta'}}{q^2}q_\mu, \\ &F^{Bq}_{1,0} = \frac{1}{\sqrt{3}}(\frac{1}{\sqrt{2}}\sin\theta F^{B\eta_8}_{1,0} + \cos\theta F^{B\eta_1}_{1,0}). \qquad (2) \end{aligned}$$

For the $\eta' - \eta$ mixing associated with decay constants above, I have used the two-angle parameterization. The numerical values of various parameters are obtained from Ref. [18] with $f_1 = 157$ MeV, $f_8 = 168$ MeV, and the mixing angles $\theta_1 = -9.1^0$, $\theta_8 = -22.1^0$. For the mixing angle associated with form factors, I used the one-angle parameterization with[18] $\theta = -15.4^o$, since these form factors were calculated in that formulation[11,7]. In the latter discussion of $b \to sg\eta'$, I shall use the same parameterization in order to compare our results with those of earlier works[10,6]. For form factors, I assume that $F^{B\eta_1} = F^{B\eta_8} = F^{B\pi}$ with dipole and monopole q^2 dependence for F_1 and F_0, respectively. I used the running mass $m_s \approx 120$ MeV at $\mu = 2.5$ GeV and $F^{B\pi} = 0.33$ following Ref.[13].

Using $V_{ts} = 0.038$, $\gamma = 64^0$ and $\mu = 5$ GeV, it is found that the branching ratios in the signal region $p_{\eta'} > 2.2$ GeV ($m_X < 2.35$ GeV) are given by

$$B(b \to \eta' s) = 0.9 \times 10^{-4}, \;\; B(B \to \eta' s\bar{q}) = 0.1 \times 10^{-4}. \tag{3}$$

The branching ratio can reach 2×10^{-4} if all parameters take values in favor of $B \to \eta' X_s$. Clearly the mechanism by four-quark operator is not sufficient to explain the observed $B \to \eta' X_s$ branching ratio.

I now turn to $b \to \eta' sg$ through the QCD anomaly. To see how the effective Hamiltonian in Eq. (1) can be applied to calculate this process, let me rearrange the effective Hamiltonian such that

$$\begin{aligned}\sum_{i=3}^{6} C_i O_i &= (C_3^{eff} + \frac{C_4^{eff}}{N_c})O_3 + (C_5^{eff} + \frac{C_6^{eff}}{N_c})O_5 \\ &-2(C_4^{eff} - C_6^{eff})O_A + 2(C_4^{eff} + C_6^{eff})O_V + C_8^{eff} O_8,\end{aligned} \tag{4}$$

where

$$O_A = \bar{s}\gamma_\mu(1-\gamma_5)T^a b \sum_q \bar{q}\gamma^\mu\gamma_5 T^a q, \;\; O_V = \bar{s}\gamma_\mu(1-\gamma_5)T^a b \sum_q \bar{q}\gamma^\mu T^a q. \tag{5}$$

Since the light-quark bilinear in O_V carries the quantum number of a gluon, one expects [10] O_V give contribution to the $b \to sg^*$ form factors. In fact, by applying the QCD equation of motion : $D_\nu G_a^{\mu\nu} = g_s \sum \bar{q}\gamma^\mu T^a q$, one obtains, $O_V = (1/g_s)\bar{s}\gamma_\mu(1-\gamma_5)T^a b D_\nu G_a^{\mu\nu}$. The effective $b \to sg^*$ vertex can be written as

$$\Gamma_\mu^{bsg} = -\frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \frac{g_s}{4\pi^2}(\Delta F_1 \bar{s}(q^2\gamma_\mu - \not{q}\, q_\mu)LT^a b - iF_2 m_b \bar{s}\sigma_{\mu\nu}q^\nu RT^a b). \tag{6}$$

The form factors ΔF_1 and F_2 are defined according to the convention in Ref. [6]. One obtains

$$\Delta F_1 = \frac{4\pi}{\alpha_s}(C_4^{eff}(\mu) + C_6^{eff}(\mu)), \;\; F_2 = -2C_8^{eff}(\mu) \tag{7}$$

Note that the relative sign of ΔF_1 and F_2 obtained agree with those in Refs. [6,7] and [19] which results in a destructive interference.

At the hadronic level the anomalous $\eta' - g - g$ coupling is given by: $a_g(\mu)\cos\theta\epsilon_{\mu\nu\alpha\beta}q^\alpha k^\beta$ with $a_g(\mu) = \sqrt{N_F}\alpha_s(\mu)/\pi f_{\eta'}$, q and k the momenta of the two gluons. Using the WC's at $\mu = 5$ GeV, the branching ratio is found to be $B(b \to sg\eta') = 5.6 \times 10^{-4}$ with a cut on $m_X \equiv \sqrt{(k+p')^2} \leq 2.35$ GeV. The spectrum for $dB(b \to sg\eta')/dm_X$ is depicted in Fig. 1. The peak of the spectrum corresponds to $m_X \approx 2.4$ GeV. The destructive interference between F_1 and F_2 lowers down the branching ratio by about 14% which is quite different from the results obtained in Refs.[6,10] because our ΔF_1 is larger than theirs because the inclusion of the matrix element corrections.



FIG. 1.

Figure 1: The distribution of $B(b \to s + g + \eta')$ as a function of the recoil mass m_X.

In the above calculation, $a_g(\mu)$ of the $\eta' - g - g$ vertex is treated as a constant independent of invariant-masses of the gluons, and μ is set to be 5 GeV. In practice, $a_g(\mu)$ should behave like a form-factor which becomes suppressed as the gluons attached to it go farther off-shell [7]. It is possible that the branching ratio obtained gets reduced significantly by the form-factor effect in $\eta' - g - g$ vertex. Should a large form-factor suppression occur, the additional contribution from $b \to \eta' s$ and $B \to \eta' s\bar{q}$ discussed earlier will become crucial. I however would like to stress that the estimate of $b \to sg\eta'$ with α_s evaluated at $\mu = 5$ GeV is conservative. To illustrate this, let me compare branching ratios for $b \to sg\eta'$ obtained at $\mu = 5$ GeV and $\mu = 2.5$ GeV respectively. The branching ratios at the above two scales with the kinematic al cut on m_X are 4.9×10^{-4} and 8.5×10^{-4} respectively. One can clearly see the significant scale-dependence! With the enhancement resulting from lowering the renormalization scale, there seems to be some room for the form-factor suppression in the attempt of explaining $B \to \eta' X_s$ by $b \to sg\eta'$.

It is clear that the Standard Model prediction for $B \to \eta' X_s$ is not in conflict with experimental data from CLEO.

2 $B \to J/\psi K^*$ and Test for CP Violation Beyond SM

The CLEO Collaboration has recently reported[5] the first full angular analysis of $B \to J/\psi K^{*0}$ decays. They find that the P wave component is small, $|P|^2 = |A_T|^2 = 0.16 \pm 0.08 \pm 0.04$, and final state interaction (FSI) phases $\phi(A_T) = -0.11 \pm 0.46 \pm 0.03\ rad$, and $\phi(A_{||}) = 3.00 \pm 0.37 \pm 0.04\ rad$ which are consistent with zero or π FSI phases in the convention where $\phi(A_0) = 0$. The small value for $|P|^2$ shows that $B \to J/\psi K^{*0} \to J/\psi K_S \pi^0$ decay is dominated by CP-even final states. This makes it practical to use $B \to J/\psi K^* \to J/\psi K_S \pi^0$ to observing mixing induced CP violation and to measure $\sin 2\beta$ without invoking an angular analysis[20]. The difference between $\sin 2\beta$ measured from $B \to J/\psi K_S$ ($\sin 2\beta_{J/\psi K_s}$) and $B \to J/\psi K^*$ ($\sin 2\beta_{J/\psi K_S \pi^0}$) can provide good test for CP violation beyond the SM[2]. With increased luminosities at CLEO, B-factories and other facilities, the charges of the B aand K^* can be identified, and therefore it is also possible to study direct CP violation in $B \to J/\psi K^*$ decays[3].

To proceed, let me show how $\sin 2\beta_{J/\psi K_S}$ and $\sin 2\beta_{J/\psi K_S \pi^0}$ can be used to test CP violation beyond the SM. The usual mixing induced CP violation measure is $\mathrm{Im}\,\xi = \mathrm{Im}\,\{(q/p)(A^* \bar{A}/|A|^2)\}$, where $q/p = e^{-2i\phi_B}$ is from B^0–$\bar{B}^0$ mixing, while A, $\bar{A}$ are B, $\bar{B}$ decay amplitudes into the same final state. For $B \to J/\psi K_S$, the final state is P-wave hence CP odd. Setting the weak phase to be σ_S, one has

$$\mathrm{Im}\,\xi(B \to J/\psi K_S) = -\sin(2\phi_B + 2\sigma_S) \equiv -\sin 2\beta_{J/\psi K_S}.$$

For $B \to J/\psi K^* \to J/\psi K_S \pi^0$, the final state has both P-amplitude (CP odd) and S- and D-amplitudes (CP even) (or their linear combinations $A_{||}$ and A_0). Let me denote their corresponding weak phases by σ_T, and $\sigma_{||}$ and σ_0, respectively. If $A_{||}$ and A_0 amplitudes have a common weak phase σ_*, one has

$$\begin{aligned} \mathrm{Im}\,\xi(B \to J/\psi K_S \pi^0) &= \mathrm{Im}\,\{e^{-2i\phi_B}[e^{-2i\sigma_T}|P|^2 - e^{-2i\sigma_*}(1 - |P|^2)]\} \\ &\equiv -(1 - 2|P|^2)\sin 2\beta_{J/\psi K_S \pi^0}, \end{aligned} \tag{8}$$

In the SM one obtains the usual result of $\sin 2\beta_{J/\psi K_S} = \sin 2\beta_{J/\psi K_S \pi^0} = \sin 2\beta$. Clearly, both measurements provide true information about $\sin 2\beta$ within SM. However, this is no longer true if one goes beyond SM. Even for the case where the weak phases of $A_{||}$ and A_0 amplitudes are equal, if $\sigma_T \neq \sigma_*$ or if $\sigma_T, \sigma_* \neq \sigma_S$, then $\sin 2\beta_{J/\psi K_S} \neq \sin 2\beta_{J/\psi K_S \pi^0}$ follows.

To the first order in new weak phases, one has

$$\Delta_{KK^*} \approx (2\sigma_S - 2\sigma_* + \frac{2|P|^2}{1 - 2|P|^2}(\sigma_T - \sigma_*))\cos(2\phi_B). \tag{9}$$

If experiments will measure a non-zero Δ_{KK^*}. It is a signal for CP violation beyond the SM.

There are many ways where new physics may change the phases σ_i. To the lowest order they may arise from dimension 6 four quark operators and dimension 5 color dipole moment operators. New physics contributions from $C_L\bar{c}\gamma^\mu(1\pm\gamma_5)c\bar{s}\gamma_\mu(1-\gamma_5)b$ type of four quark interactions are proportional to the dominant SM contribution in Eq.[1] which just generate a common weak phase for all the amplitudes and therefore Δ_{KK^*} is zero like the SM prediction. Δ_{KK^*} discussed here does not provide good test for new physics of this type. The interaction of the form $C_R\bar{c}\gamma^\mu(1\pm\gamma_5)c\bar{s}\gamma_\mu(1+\gamma_5)b$ will generate different phases for A_T and $A_{||,0}$. The reason is that the current $\bar{s}\gamma^\mu(1+\gamma_5)b$ contribution to A_T is proportional to $C_{SM}+C_R$, but to $A_{||,0}$ is proportional to $C_{SM}-C_R$. To a good approximation, $\delta_T=-\delta_{||,0}$. Δ_{KK^*} can be different from zero.

As an example, let us study R-parity violating supersymmetric models. In R-parity violating supersymmetric models, exchange of charged sleptons or down type squarks can generate non-zero C_R with an arbitrary phase δ_R. The allowed value for C_R is constrained from experimental data on $b\to s\gamma$. This still allow C_R term to contribute to $B\to J/\psi K^*$ at the amplitude level as large as 20% of the SM contribution. Stronger constraints can be obtained by assuming that $b\to c\bar{c}s$ is similar in strength to $b\to c\bar{u}s$ [21]. The upper bound of the weak phases are approximately given by

$$\delta_T=-\delta_*\approx 0.1\sin\delta_R. \tag{10}$$

Using Eq.(9), one can easily see that the difference Δ_{KK^*} can reach $-0.35\sin\delta_R\cos(2\phi_B)$. Such a large difference can be detected at B factories.

The contribution from dimension 5 color dipole operator has been estimated by assuming that color octet operators contribute the amount as determined in generalized factorization approximation. The magnitude of the color dipole coefficient as large as 10 times of the SM, if the chiral structure is the same as the SM one, is not ruled out and may in fact play some important role in the missing charm problem. The weak phases $\sigma_{T,||,0}$ can be as large as $0.08\sin\delta_c$, where δ_c is the weak phase of the color dipole phase [2]. The phases are approximately equal for σ_T and σ_*. The phase σ_S is suppressed by a factor of m_ψ^2/m_B^2 due to helicity structure of the operator. The value for Δ_{KK^*} can be as large as $0.18\sin\delta_c\cos(2\phi_B)$. If it turns out that the color dipole interaction has opposite chirality, although there is an enhancement factor of $1/(1-2|P|^2)$, the coefficient C_R is more stringently constrained resulting in a smaller Δ_{KK^*}.

I now discuss direct CP violation in angular distribution for charge $B \to J/\psi K^*$. The full angular distribution for $B \to J/\psi K^*$ is given by

$$\begin{aligned}
&\frac{1}{\Gamma}\frac{d\Gamma}{d\cos\theta_{tr} d\cos\theta_{K^*} d\phi_{tr}} = \frac{9}{32\pi}\{2|A_0|^2\cos^2\theta_{K^*}(1-\sin^2\theta_{tr}\cos^2\phi_{tr}) \\
&+|A_{||}|^2\sin^2\theta_{K^*}(1-\sin^2\theta_{tr}\sin^2\phi_{tr}) + |A_T|^2\sin^2\theta_{K^*}\sin^2\phi_{tr} \\
&-Im(A_{||}^* A_T)\sin^2\theta_{K^*}\sin 2\theta_{tr}\sin\phi_{tr} \\
&+\frac{1}{\sqrt{2}}Re(A_0^* A_{||})\sin 2\theta_{K^*}\sin^2\theta_{tr}\sin 2\phi_{tr} \\
&+\frac{1}{\sqrt{2}}Im(A_0^* A_T)\sin 2\theta_{K^*}\sin 2\theta_{tr}\cos\phi_{tr}\}, \qquad (11)
\end{aligned}$$

where the transversity angles θ_{tr} and ϕ_{tr} are defined as polar and azimuth angles of the charge lepton in the J/ψ rest frame with x axis along the direction of K^*, x-y plane parallel to $K\pi$ plane. The angle θ_{K^*} is defined as that of the K in the rest frame of K^* relative to the negative of the J/ψ direction in that frame.

In the CLEO analysis, the FSI phase for A_0 was taken to be zero. For convenience I will use the convention that each amplitude A_i has both CP conserving FSI phase ϕ_i and CP violating phase σ_i as indicated before. With this convention, $A_j = |A_j|e^{i(\phi_j+\sigma_j)}$ while $\bar{A}_T = -|A_T|e^{i(\phi_T-\sigma_T)}$, $\bar{A}_{(||,0)} = |A_{(||,0)}|e^{i(\phi_{(||,0)}-\sigma_{(||,0)})}$.

It is clear that the coefficients $\alpha = -Im(A_{||}^* A_T)$, $\beta = Re(A_0^* A_{||})$, and $\gamma = Im(A_0^* A_T)$ of the last three terms in the angular distribution, and similarly $\bar{\alpha}$, $\bar{\beta}$ and $\bar{\gamma}$ for $\bar{B}$ decays, contain information about CP violation. Without separating B and $\bar{B}$ decays, however, which was the case for the CLEO analysis mentioned earlier, information on CP violation cannot be extracted. One must obtain the angular distributions for $B \to J/\psi K^*$ and $\bar{B} \to J/\psi \bar{K}^*$ decays separately and determine the coefficients for the interference terms in each case. The following three quantities then measure CP violation [3,22]

$$\begin{aligned}
a_1 &= \alpha + \bar{\alpha} = 2|A_{||}||A_T|\cos(\phi_{||T})\sin(\sigma_{||T}), \\
a_2 &= \beta - \bar{\beta} = -2|A_{||}||A_0|\sin(\phi_{||0})\sin(\sigma_{||0}), \\
a_3 &= \gamma + \bar{\gamma} = -2|A_T||A_0|\cos(\phi_{T0})\sin(\sigma_{T0}), \qquad (12)
\end{aligned}$$

where $\phi_{ij} = \phi_i - \phi_j$ and $\sigma_{ij} = \sigma_i - \sigma_j$.

It is interesting to note that the CP violating observable $a_{1,3}$ do not require FSI phase differences and is especially sensitive to CP violating weak phases.

The present CLEO data on angular distributions which provide information about FSI phases are the charge averaged CP conserving quantities which are proportional to $\sin(\phi_{||T})\cos(\sigma_{||T})$, $\cos(\phi_{||0})\cos(\sigma_{||0})$, and $\sin(\phi_{T0})\cos(\sigma_{T0})$. At present these data do not exclude $a_{1,3}$ up to 50% and smaller for a_2

Again in the SM, the three CP violating observables are zero. They provide good tests for CP violation beyond the SM there it is possible to have non-zero values for a_i. For example in R-parity violating supersymmetric model considered earlier, for the central values for the magnitude of the amplitudes and assuming that the FSI are zero, the asymmetries can be as large as $a_1 = 0.10\sin\delta_R$ and $a_3 = -0.12\sin\delta_R$.

The color dipole interaction to these asymmetries are small because with the same chiral structure as the SM, the new phases are approximately equal as discussed before and therefore small asymmetries. For the case with opposite chiral structure as the SM the strength of the interaction is more stringently constrained, the phases are all small.

In all cases discussed above the weak phases for $\sigma_{||}$ and σ_0 are equal (or approximately equal). The asymmetry a_2 is approximately zero in all cases considered, and does not seem to be a good quantity to study for CP violation using this method.

The sensitivities for $a_{1,2}$ is similar to the sensitivity to the phase angles of the amplitudes. It is interesting to note that the systematic error in CLEO analysis is already as low as 0.03 [5]. With increases statistics, $a_{1,3}$ as large as 0.10 should be accessible at CLEO, at CDF, and at future B factories. Since the errors are determined through a fit, it is not clear how the statistical error scales with actual number of events. The question can only be answered by actual studies, but naive scaling implies that one would need 10^8 events to be able to distinguish the deviations given earlier. Nevertheless, the needed number of events may be less and measurement of the observables a_i will provide us with useful information about CP violation.

I thank W.-S. Hou and G.-L. Lin for collaborations on the work reported here. This work is supported in part by ROC National Science Council and by Australian Research Council.

1. Xiao-Gang He and Guey-Lin Lin, e-print hep-ph/9809204.
2. Xiao-Gang He and Wei-Shu Hou, e-print hep-ph/9804293.
3. Xiao-Gang He and Wei-Shu Hou, e-print hep-ph/9807507 (Phys. Rev. D in press).
4. CLEO Collaboration T. E. Browder *et al.*, Phys. Rev. Lett. 81, 1786(1998).
5. CLEO Collaboration C.P. Jessop et al., Phys. Rev. Lett. **79**, 4533 (1997).

6. W.S. Hou and B. Tseng, Phys. Rev. Lett. **80**, 434 (1998).
7. A. L. Kagan and A. Petrov, e-print hep-ph/9707354; A. L. Kagan, hep-ph/98-6266.
8. H. Fritzsch, Phys. Lett. **B 415**, 83 (1997); X.-G. He, W.-S. Hou and C.S. Huang, Phys. Lett. **B429**, 99 (1998).
9. CLEO collaboration, B.H. Behrens et al., Phys. Rev. Lett. **80**, 3710 (1998).
10. D. Atwood and A. Soni, Phys. Lett. **B 405**, 150 (1997).
11. A. Datta, X.-G. He, and S. Pakvasa, Phys. Lett. **B 419**, 369 (1998).
12. H.-Y. Cheng, and B. Tseng, e-print hep-ph/9803457; A. Ali, J. Chay, C. Greub and P. Ko, Phys. Lett. **B424**, 161 (1998); N. Deshpande, B. Dutta and S. Oh, Phys. Rev. **D57**, 5723(1998).
13. A. Ali, G. Kramer and C.-D. Lu, hep-ph/9804363.
14. I. Halperin and A. Zhitnitsky, Phys. Rev. Lett. **80**, 438 (1998); F. Araki, M. Musakonov and H. Toki, e-print hep-ph/9803356; D.S. Du, Y.-D. Yang and G.-H. Zhu, e-print hep-ph/9805451; M. Ahmady, E. Kou and A. Sugamoto, Phys. Rev. **D58**, 014015 (1998); D.S. Du, C.S. Kim and Y.-D. Yang, Phys. Lett. **B426**, 133 (1998); F. Yuan and K.-T. Chao, Phys. Rev. **D 56**, 2495 (1997); A. Dighe, M. Gronau and J. Rosner, Phys. Rev. Lett. **79**, 4333 (1997).
15. For a review see G. Buchalla, A. J. Buras and M. E. Lautenbacher, Review of Modern Physics, **68**, 1125 (1996).
16. N. G. Deshpande and X.-G. He, Phys. Lett. **B 336**, 471 (1994).
17. M. Ciuchini et al., Nucl. Phys. **B421**, 41(1994).
18. T. Feldmann, P. Kroll and B. Stech, hep-ph/9802409.
19. H. Simma and D. Wyler, Nucl. Phys. **B344**, 283(1990); A. Lenz, N. Nierste and G. Ostermier, Phys. Rev. **D56**, 7228(1997).
20. I. Dunietz et al., Phys. Rev. D**43**, 2193 (1991); A.S. Dighe et al., Phys. Lett. B**369**, 144 (1996).
21. Y. Grossman and M. Worah, Phys. Lett. B**395**, 241 (1997).
22. German Valencia, Phys. Rev. **D39**, 3339(1989).